



FMUP FACULDADE DE MEDICINA
UNIVERSIDADE DO PORTO

MESTRADO INTEGRADO EM MEDICINA

2017/2018

Ema Catarina Monteiro dos Santos

**Fluorescence-guided resection with 5-aminolevulinic acid:
implications, complications and learning after 100 cases**

março, 2018

FMUP

Ema Catarina Monteiro dos Santos

Fluorescence-guided resection with 5-aminolevulinic acid:
implications, complications, and learning after 100 cases

Resseção guiada por fluorescência com ácido 5-aminolevulinico:
implicações, complicações e aprendizagem após 100 casos

Mestrado Integrado em Medicina

Área: Neurocirurgia

Tipologia: Dissertação

Trabalho efetuado sob a Orientação de:

Paulo José Campos Linhares Vieira

Trabalho organizado de acordo com as normas da revista:

Acta Neurochirurgica

março, 2018

Eu, **Ema Catarina Monteiro dos Santos**, abaixo assinado, nº mecanográfico **201202274**, estudante do 6º ano do Ciclo de Estudos Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 19/03/2018

Assinatura conforme cartão de identificação:

Ema Catarina Monteiro dos Santos

NOME

Ema Catarina Monteiro dos Santos

NÚMERO DE ESTUDANTE

201202274

E-MAIL

emamsantos@gmail.com

DESIGNAÇÃO DA ÁREA DO PROJECTO

Neurocirurgia

TÍTULO DISSERTAÇÃO

Fluorescence-guided resection with 5-aminolevulinic acid: implications, complications, and learning after 100 cases

ORIENTADOR

Paulo José Campos Linhares Vieira

COORDINADOR (se aplicável)

-

ASSINALE APENAS UMA DAS OPÇÕES:

É AUTORIZADA A REPRODUÇÃO INTEGRAL DESTA OBRA APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input type="checkbox"/>
É AUTORIZADA A REPRODUÇÃO PARCIAL DESTA OBRA (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input type="checkbox"/>
DE ACORDO COM A LEGISLAÇÃO EM VIGOR, (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) NÃO É PERMITIDA A REPRODUÇÃO DE QUALQUER PARTE DESTA OBRA.	<input checked="" type="checkbox"/>

Faculdade de Medicina da Universidade do Porto, 19/03/2018

Assinatura conforme cartão de identificação: Ema Catarina Monteiro dos Santos

Fluorescence-guided resection with 5-aminolevulinic acid: implications, complications and learning after 100 cases

Ema Santos¹, Bruno Carvalho^{1,2}, Rui Vaz,^{1,2,3} Paulo Linhares^{1,2,3}

¹ Faculty of Medicine, Oporto University, Oporto, Portugal

² Department of Neurosurgery, Hospital São João, São João Hospital Centre, Oporto, Portugal

³ Neurosciences Center, Hospital CUF Porto, Oporto, Portugal

Corresponding author:

Ema Catarina Monteiro dos Santos

Faculty of Medicine, Oporto University, Oporto, Portugal

Alameda Prof. Hernâni Monteiro, 4200-319 Porto

E-mail: emamsantos@gmail.com

Abstract

Background: 5-aminolevulinic acid (5-ALA) has demonstrated to be a safe and useful tool for real-time intraoperative identification of tumor tissue, leading to extensive resections which are associated with increased survival for patients with high-grade gliomas. Although tumors in eloquent areas can constrain resection due to higher risk of neurological deficits, this should not contraindicate 5-ALA judicious use.

Objective: To evaluate the impact of resection in progression-free survival (PFS) and overall survival (OS), the postoperative complications and learning after the first 100 cases of 5-ALA fluorescence-guided resection in a tertiary referral hospital.

Methods: A retrospective study of the first 100 cases of newly diagnosed brain lesions submitted to fluorescence-guided surgery with 5-ALA, from 2012 to 2016. Only glioblastoma patients were considered for the statistical analysis. We analyzed the influence of 5-ALA in the extent of resection considering tumor in eloquent and non-eloquent location, post-operative neurological deficits, PFS and OS.

Results: Gross total resection (GTR) was achieved in 73% of patients with tumors in non-eloquent areas and in 43,4% in eloquent areas. Partial resection was superior in tumors located in eloquent areas (34,0%vs13,5%). There were no significant differences in postoperative neurological deficits amongst distinct extents of resection ($p=0,228$) or tumor location ($p=0,670$). Mean PFS for all group was $9,73\pm1,19$ months and mean OS for all group was $16,27\pm1,39$ months. $GTR>90\%$ and the non-eloquent area of the tumor were correlated with a significant increase in PFS but the extent of resection was the only independent factor for PFS in multivariate analysis. Regarding OS, no independent prognosis factor was established in the multivariate analysis.

Conclusion: 5-ALA is an important tool in glioblastoma surgery, contributing for GTR and, consequently, to increase PFS. Although its use has not raised neurological deficits, it must be carefully used in eloquent located tumors.

Key-words: 5-aminolevulinic acid; fluorescence-guided resection; glioblastoma; extent of resection;

Introduction

The extent of resection has been recognized as an established prognostic factor in the treatment of glioblastoma (GBM) [13,1], being the only one potentially modifiable. Complete resection of enhancing tumor is linked to an improvement in survival [2,17,16,13,21], acting synergistically with optimized medical treatment [18].

Multiple techniques were developed in order to achieve a better identification of the tumor from adjacent normal brain, being fluorescence-guided resection with 5-aminolevulinic acid (5-ALA) one of the most innovative.

5-ALA is a metabolic precursor in the heme pathway and can be used as an oral prodrug that leads to the synthesis and accumulation of fluorescent porphyrins, mainly protoporphyrin IX, in high-grade glioma cells [20,19,15].

Several studies have showed that 5-ALA intraoperative fluorescence has a very high sensitivity, specificity and positive predictive value (PPV) for malignant tumor tissue [22,15,7,26,23,11,8,10]. Additionally, there are two patterns of fluorescence: “strong”, associated with solid and proliferating tumor and “weak” corresponding to infiltrating tumor with both tumor and normal cells. These patterns correlate with sensitivity, specificity and PPV, and has been found that these values can reach 100% in areas with “strong” fluorescence [23,7,11,12,10], confirming the remarkable diagnostic precision of 5-ALA.

Introduced by Stummer et al.[22], it has since demonstrated to be a secure and accurate tool for real-time intraoperative identification of tumor cells, leading to extensive tumor removal and an increase in survival for patients with high-grade gliomas[20,19,5]. It remains inconsistent in the literature if the increased survival expresses progression-free survival or overall survival.

However, it cannot be ignored that more radical resections carry more risks of neurological damages [24], particularly in tumors near eloquent areas. The aim of using 5-ALA in eloquent areas is not to achieve gross total resection (GTR) but increase the accuracy for tumor cells. While some authors only use immunofluorescence in cases of expected GTR, our understanding is that 5-ALA also helps in defining tumor cells even in cases only suitable for partial resections (PR).

The goals of the present study are to evaluate the impact of the extent of tumor resection in progression-free survival and overall survival, the possible

post-surgical complications and the lessons learned after the first 100 cases using fluorescence-guided resection with 5-ALA.

Materials and Methods

Patient population

The data used in this study was obtained from clinical records of the first 100 cases submitted to fluorescence-guided surgery with 5-ALA, from May 2012 until February 2016 in a European Union University Hospital. All medical record reviews were conducted with approval from the Local Ethical Committee. For this type of study formal consent is not required.

The criteria for the use of 5-ALA were newly diagnosed brain lesions with MRI suggestive of high grade malignant glioma in patients eligible for open surgery, according to their clinical status.

Tumors were categorized as located in eloquent and non-eloquent areas. The primary motor cortex (precentral gyrus), primary somatosensory cortex (postcentral gyrus), occipital visual cortex, insula, corpus callosum and the operculum area in the frontal, parietal and temporal left lobes were considered eloquent areas.

From a total of 100 patients submitted to surgery with 5-ALA, 90 were diagnosed with GBM after histopathological confirmation. In the remaining 10 patients, 3 were diagnosed with anaplastic astrocytoma and 7 with different histological types. Due to the diversity of diagnoses, patients who were not diagnosed with GBM were excluded in the further analysis. **Table 1** shows the patient and tumor baseline characteristics.

Technical principals

5-ALA (Gliolan, Medac, Wedel, Germany) oral solution was administered to patients at 20mg/kg bodyweight, 2-4h before general anesthesia. Dexamethasone was also provided to all patients at least two days before surgery up to the postoperative MRI, less than 72h after surgery. It has been reported that dexamethasone use improves fluorescence.

The tumor was resected using a surgical microscope (*Zeiss OPMI Pentero 900®* or *Leica OHS®*) that is able to switch from white conventional light (no fluorescence visible) to violet-blue light (BL400) whenever desired by the

neurosurgeon, allowing fluorescent tumor tissue identification. The goal of the surgery was to remove safely all the fluorescent tissue visible, considering possible neurological deterioration when near to eloquent areas. Other intraoperative diagnostic tools, such as neuronavigation and awake monitoring were used.

The extent of resection was assessed by comparison of preoperative MRI with postoperative MRI less than 72h after surgery and then classified as GTR, subtotal resection (STR) or PR. GTR was defined as absence of residual contrast in post-op MRI; the resection was considered subtotal when was superior to 90% of the volume but not complete; PR included less than 90% of the tumor volume removed.

Treatment and follow up

After diagnostic confirmation of GBM, patients were referred to a neuro-oncology multidisciplinary team (MDT) meeting to evaluate their clinical conditions to pursue further treatment.

From the patients diagnosed with GBM, 76 started the Stupp protocol, radiotherapy with concomitant and adjuvant temozolomide[25], one patient with recurrent brain lesion started second-line chemotherapy after surgery and six patients were referred to palliative care. Seven patients were not submitted to any treatment after surgery due to their condition (n=6) or treatment refusal (n=1) and were submitted to best supportive care.

Progression was defined as an increase in residual tumor volume, the occurrence of a new tumor lesion or deterioration in the clinical status, according to RANO criteria. After progression, the neuro-oncology MDT would decide whether to start second-line chemotherapy, re-operate or refer to palliative chemotherapy according to the clinical and functional condition of each patient.

Statistical analysis

The statistical analysis was performed using IBM SPSS® (Statistical Package for Social Sciences) version 24. Statistical significance was set at $p < 0,05$. It was conducted a descriptive analysis of all variables.

To test the possible effect of multiple factors like age, sex, preoperative KPS and area of the tumor (eloquent/non-eloquent) in the extent of resection, separately Chi-square tests were performed. It was also assessed the influence of the resection on post-operative neurological deficits with a Chi-square test.

Progression-free survival and overall survival were defined as the time between surgery and progression or time of death, respectively. To identify factors influencing survival, Kaplan-Meier survival curves were constructed individually for sex, age, preoperative KPS, tumor eloquence and extent of resection. For survival analysis, the extent of resection was grouped in gross total/subtotal (tumor removal superior to 90%) and partial resection due to the quantitative difference of tumor not removed.

It was also analyzed if the extent of resection suffered a variation through with time, comparing the extent of resection of the first 45 patients with GBM submitted to 5-ALA guided-resection with the last 45 patients.

Results

Resection

Patients were divided in two groups: tumors in non-eloquent areas (n=37) and tumors in eloquent areas (n=53). It was observed a significant difference in the extent of resection between the groups ($p=0,019$) (FIG1). GTR was accomplished in 73% of patients with non-eloquent tumors, 13,5% had STR and 13,5% PR, which means that it was obtained a resection >90% in 86,5% of patients. Concerning tumors in eloquent areas, resections >90% were achieved in 66% being GTR 43,4% of the patients, 22,6% had STR of the tumor and 34,0% had PR. Even though in both groups GTR was achieved more often than subtotal or partial, these results demonstrate a bigger proportion of gross total removal in tumors in non-eloquent areas and a higher percentage of PR in eloquent areas, what was expected since the aim of resection in eloquent areas were the maximal safe resection and not GTR.

Factors which could have influenced the extent of resection such as age, sex, preoperative KPS and time within series (Table 2) were also considered. The analysis demonstrated that sex, age and preoperative KPS could not prove to be significant prognosis factors even though it appears that a GTR is accomplished more frequently in younger patients or with KPS score superior to

70. The extent of resection did not suffer a variation when comparing the first 45 patients submitted to 5-ALA guided resection with the latter 45 patients.

Safety

Post-operative complications were divided in de novo epilepsy (n=2), hemorrhagic/ ischemic events (n=3), infection (n=2) and new/aggravation of focal deficit (n=20). The majority of patients (70%) have not developed any complication. KPS score 1 month after surgery remained identical to pre-op KPS score (median=90).

From all the post-operative complications detected, it was considered that only new/aggravation of focal deficit could result from the use of 5-ALA since its main goal is to improve the extent of resection, which can ultimately lead to neurological deterioration. It was also considered if focal deficits could vary with the area of the tumor (eloquent/non-eloquent) as eloquent areas would be more prone to subsequent deficits.

After analysis, no difference in post-op focal deficit was found amongst tumors in eloquent areas comparing with non-eloquent areas ($p=0,670$). Moreover, the neurological deficits have not diverged when compared between gross total, subtotal or partial resection ($p=0,228$).

Survival

Mean overall survival for all group was $16,27 \pm 1,39$ months and mean progression-free survival for all group was $9,73 \pm 1,19$ months.

For the analysis of survival, potential prognosis factors were evaluated such as sex, patient age, preoperative KPS score, localization of tumor and the extent of resection.

Patient age inferior to 65 years old ($<65 = 18,01 \pm 1,57$ months vs $\geq 65 = 13,22 \pm 2,31$ months; $p=0,043$) and a preoperative KPS score superior to 70 ($KPS > 70 = 17,36 \pm 1,54$ months vs $KPS \leq 70 = 10,13 \pm 2,00$ months; $p=0,028$) were significantly associated with longer overall survival. Extent of resection ($GTR > 90\% = 17,52 \pm 1,68$ months vs $PR = 13,130 \pm 2,62$ months; $p=0,107$) and tumor area (eloquent= $13,71 \pm 1,32$ months vs non-eloquent= $19,32 \pm 2,41$ months; $p=0,077$) failed to demonstrate a significant association, even though it appears to exist a tendency for improvement of overall survival when the resection is

>90% or the tumor is located in a non-eloquent area (**FIG. 2**). No independent prognosis factor for overall survival was established in the multivariate analysis.

Furthermore, a complete resection (GTR>90%=11,33±1,54 months vs PR=5,30±0,93 months; p=0,001), the non-eloquent area of the tumor (eloquent=7,15±0,81 months vs non-eloquent=13,58±2,59 months; p=0,007) and once again, the preoperative KPS score superior to 70 (KPS>70=10,48±1,37 months vs KPS≤70=5,49±1,10 months; p=0,028) showed a significant increase in progression-free survival. The difference in patient age was not statistically significant in progression-free survival (<65=9,78±1,17 months vs ≥65=9,22±2,25 months; p=0,308) (**FIG 3**). Multivariate analysis was performed, showing that resection superior to 90% is the only independent factor for progression-free survival (p=0,030)

There were no differences in overall survival (male=17,07±2,03 months vs female=14,83±1,57months; p=0,590) or progression-free survival (male=9,66±1,50 months vs female=9,19±1,51months; p=0,820) related with gender.

Discussion

Glioblastoma is the most common malignant primary brain tumor and it is associated with a very poor prognosis. Tumor removal has a major role in treatment, being proven that the extent of resection is an important prognosis factor for patient survival [21,17,16,13,2,1]. This led to the development of 5-ALA with the goal of achieving more radical resections thus improving survival. [22,21,20]

Multiple studies revealed GTR rates superior to 60% using 5-ALA, representing a major improvement from “white-light” surgery [20,19,8,7,3]. Although 5-ALA is used mostly in patients where the goal is GTR, we used it in all patients with glioblastoma regardless of the expected extent of resection. Because of this, we divided our results in two different groups, the one that is expected a GTR and the group which STR was the aim because of an eloquent area is present. Tumors in eloquent areas have a particular importance due to increased risk of neurological damage and can substantially limit the resection. One must not forget the importance of the quality of life in patients with such a short life expectancy and improving resection does not fulfill its goal if

accompanied with new or aggravation of neurological deficits and can even decrease overall survival.[14] These leads to differences in the extent of resection when comparing tumors in eloquent and non-eloquent areas.

In the present study, resection >90% of tumor volume was observed in 86,5% and GTR in 73% of patients with tumors located in non-eloquent areas, in contrast with only 66% with resection >90% and 43,4% with complete absence of contrast of tumors located near an eloquent area. On the other hand, the proportion of PR and STR was superior in tumors adjacent to eloquent areas. These results reflect a careful approach to the use of 5-ALA specially in eloquent areas, not removing all the fluorescent tissue that is observed intraoperatively to prevent further neurological damages. Other intraoperative diagnostic tools were used, but in order to achieve better results in resection it has been suggested the use of intraoperative monitoring in combination with 5-ALA in all tumors located in eloquent areas. [9,6]

It was detected that the extent of resection did not suffer a modification when comparing the first 45 patients submitted to 5-ALA guided resection with the latter 45 patients. This lack of variation through time can be rationalized by the fact that few surgeons started performing this type of surgery progressively and their diverse learning curves.

Regarding safety, some authors described a temporarily neurological impairment in patients with more radical resections[24]. We have not experienced differences in postoperative focal deficit amongst tumors in eloquent areas comparing with non-eloquent areas or when compared between GTR, STR or PR, although this can be consequence of the careful approach and higher proportion of PR and STR in eloquent tumors.

Our mean overall survival for all group was $16,27 \pm 1,39$ months and mean progression-free survival for all group was $9,73 \pm 1,19$ months which is concordant with the results in the literature [11] even with lower resection rates.

Exploring overall survival, we observed that younger ages and a good preoperative status were significantly associated with longer overall survival in univariate analysis but not in multivariate analysis. Additionally, it appeared to exist a non-significant tendency for improvement of overall survival when the tumor is located in a non-eloquent area or the resection is >90%, what would be consistent with previous results referring a survival benefit in resections superior

to 70% [17,2]. In spite of that, OS could be a controversial end-point owing to the difficulty to control treatments and interventions after progression. In fact, Bloch et al.[4] stated that the extent of resection at recurrence can be an essential predictor of overall survival. The differences in post-recurrence treatments or re-operations were outside the sphere of this work but could explain the lack of statistical significance in respect of resection and OS.

A significant improvement in progression-free survival was noted with GTR>90%, the non-eloquent area of the tumor and a favorable preoperative status. In multivariate analysis, resection superior to 90% was the only independent factor. PFS was also the clinical benefit described by Stummer et al.[20]

Concerning limitations, the retrospective design of the study and the small sample size are important factors and can explain the lack of statistical significance in some variables. Further prospective studies with larger samples are needed to help recognize statistical significance in all the analysis a tendency was acknowledged and to improve the evidence level for the use of 5-ALA guided resection.

Conclusions:

5-ALA is an important tool in glioblastoma surgery, contributing for GTR and, consequently, to duplicate PFS from 5,30 to 11,33 months. Although its use has not raised neurological deficits, it must be carefully used in eloquent tumors. Our data were according to the literature, showing better results when 5-ALA were used in tumors in non-eloquent areas where the aim is to achieve gross total resection.

Conflict of Interest:

The authors declare that they have no conflict of interest.

1. Chaichana K, Parker S, Olivi A, Quinones-Hinojosa A (2010) A proposed classification system that projects outcomes based on preoperative variables for adult patients with glioblastoma multiforme. *J Neurosurg* 112:997-1004. doi:10.3171/2009.9.jns09805
2. Chaichana KL, Jusue-Torres I, Navarro-Ramirez R, Raza SM, Pascual-Gallego M, Ibrahim A, Hernandez-Hermann M, Gomez L, Ye X, Weingart JD, Olivi A, Blakeley J, Gallia GL, Lim M, Brem H, Quinones-Hinojosa A (2014) Establishing percent resection and residual volume thresholds affecting survival and recurrence for patients with newly diagnosed intracranial glioblastoma. *Neuro Oncol* 16:113-122. doi:10.1093/neuonc/not137
3. Coburger J, Hagel V, Wirtz CR, Konig R (2015) Surgery for Glioblastoma: Impact of the Combined Use of 5-Aminolevulinic Acid and Intraoperative MRI on Extent of Resection and Survival. *PLoS One* 10:e0131872. doi:10.1371/journal.pone.0131872
4. Coburger J, Wirtz CR, Konig RW (2017) Impact of extent of resection and recurrent surgery on clinical outcome and overall survival in a consecutive series of 170 patients for glioblastoma in intraoperative high field magnetic resonance imaging. *Journal of neurosurgical sciences* 61:233-244. doi:10.23736/s0390-5616.16.03284-7
5. Cordova JS, Gurbani SS, Holder CA, Olson JJ, Schreibmann E, Shi R, Guo Y, Shu HK, Shim H, Hadjipanayis CG (2016) Semi-Automated Volumetric and Morphological Assessment of Glioblastoma Resection with Fluorescence-Guided Surgery. *Mol Imaging Biol* 18:454-462. doi:10.1007/s11307-015-0900-2
6. Della Puppa A, De Pellegrin S, d'Avella E, Gioffre G, Rossetto M, Gerardi A, Lombardi G, Manara R, Munari M, Saladini M, Scienza R (2013) 5-aminolevulinic acid (5-ALA) fluorescence guided surgery of high-grade gliomas in eloquent areas assisted by functional mapping. Our experience and review of the literature. *Acta Neurochir (Wien)* 155:965-972; discussion 972. doi:10.1007/s00701-013-1660-x
7. Díez Valle R, Tejada Solis S, Idoate Gastearena MA, García de Eulate R, Domínguez Echávarri P, Aristu Mendiroz J (2011) Surgery guided by 5-aminolevulinic fluorescence in glioblastoma: volumetric analysis of extent of resection in single-center experience. *Journal of Neuro-Oncology* 102:105-113. doi:10.1007/s11060-010-0296-4
8. Eljamel S (2015) 5-ALA Fluorescence Image Guided Resection of Glioblastoma Multiforme: A Meta-Analysis of the Literature. *Int J Mol Sci* 16:10443-10456. doi:10.3390/ijms160510443
9. Feigl GC, Ritz R, Moraes M, Klein J, Ramina K, Gharabaghi A, Krischek B, Danz S, Bornemann A, Liebsch M, Tatagiba MS (2010) Resection of malignant brain tumors in eloquent cortical areas: a new multimodal approach combining 5-aminolevulinic acid and intraoperative monitoring. *J Neurosurg* 113:352-357. doi:10.3171/2009.10.JNS09447
10. Hadjipanayis CG, Widhalm G, Stummer W (2015) What is the Surgical Benefit of Utilizing 5-Aminolevulinic Acid for Fluorescence-Guided Surgery of Malignant Gliomas? *Neurosurgery* 77:663-673. doi:10.1227/NEU.0000000000000929
11. Halani SH, Adamson DC (2016) Clinical utility of 5-aminolevulinic acid HCl to better visualize and more completely remove gliomas. *Onco Targets Ther* 9:5629-5642. doi:10.2147/OTT.S97030

12. Hefti M, von Campe G, Moschopoulos M, Siegner A, Looser H, Landolt H (2008) 5-aminolevulinic acid induced protoporphyrin IX fluorescence in high-grade glioma surgery: a one-year experience at a single institution. *Swiss medical weekly* 138:180-185. doi:2008/11/smw-12077
13. Lacroix M, Abi-Said D, Fourney DR, Gokaslan ZL, Shi W, DeMonte F, Lang FF, McCutcheon IE, Hassenbusch SJ, Holland E, Hess K, Michael C, Miller D, Sawaya R (2001) A multivariate analysis of 416 patients with glioblastoma multiforme: prognosis, extent of resection, and survival. *J Neurosurg* 95:190-198. doi:10.3171/jns.2001.95.2.0190
14. McGirt MJ, Mukherjee D, Chaichana KL, Than KD, Weingart JD, Quinones-Hinojosa A (2009) Association of Surgically Acquired Motor and Language Deficits on Overall Survival after Resection of Glioblastoma Multiforme. *Neurosurgery* 65:463-470. doi:10.1227/01.NEU.0000349763.42238.E9
15. Nabavi A, Thurm H, Zountsas B, Pietsch T, Lanfermann H, Pichlmeier U, Mehdorn M, Group ALARGS (2009) Five-aminolevulinic acid for fluorescence-guided resection of recurrent malignant gliomas: a phase ii study. *Neurosurgery* 65:1070-1076; discussion 1076-1077. doi:10.1227/01.NEU.0000360128.03597.C7
16. Sanai N, Berger MS (2008) Glioma extent of resection and its impact on patient outcome. *Neurosurgery* 62:753-764; discussion 764-756. doi:10.1227/01.neu.0000318159.21731.cf
17. Sanai N, Polley MY, McDermott MW, Parsa AT, Berger MS (2011) An extent of resection threshold for newly diagnosed glioblastomas. *J Neurosurg* 115:3-8. doi:10.3171/2011.2.jns10998
- 10.3171/2011.7.jns10238
18. Stummer W, Meinel T, Ewelt C, Martus P, Jakobs O, Felsberg J, Reifenberger G (2012) Prospective cohort study of radiotherapy with concomitant and adjuvant temozolomide chemotherapy for glioblastoma patients with no or minimal residual enhancing tumor load after surgery. *J Neurooncol* 108:89-97. doi:10.1007/s11060-012-0798-3
19. Stummer W, Novotny A, Stepp H, Goetz C, Bise K, Reulen HJ (2000) Fluorescence-guided resection of glioblastoma multiforme by using 5-aminolevulinic acid-induced porphyrins: a prospective study in 52 consecutive patients. *J Neurosurg* 93:1003-1013. doi:10.3171/jns.2000.93.6.1003
20. Stummer W, Pichlmeier U, Meinel T, Wiestler OD, Zanella F, Reulen H-J (2006) Fluorescence-guided surgery with 5-aminolevulinic acid for resection of malignant glioma: a randomised controlled multicentre phase III trial. *The Lancet Oncology* 7:392-401. doi:10.1016/s1470-2045(06)70665-9
21. Stummer W, Reulen HJ, Meinel T, Pichlmeier U, Schumacher W, Tonn JC, Rohde V, Oppel F, Turowski B, Woiciechowsky C, Franz K, Pietsch T (2008) Extent of resection and survival in glioblastoma multiforme: identification of and adjustment for bias. *Neurosurgery* 62:564-576; discussion 564-576. doi:10.1227/01.neu.0000317304.31579.17
22. Stummer W, Stocker S, Wagner S, Stepp H, Fritsch C, Goetz C, Goetz AE, Kiefmann R, Reulen HJ (1998) Intraoperative detection of malignant gliomas by 5-aminolevulinic acid-induced porphyrin fluorescence. *Neurosurgery* 42:518-525; discussion 525-516

23. Stummer W, Tonn JC, Goetz C, Ullrich W, Stepp H, Bink A, Pietsch T, Pichlmeier U (2014) 5-Aminolevulinic acid-derived tumor fluorescence: the diagnostic accuracy of visible fluorescence qualities as corroborated by spectrometry and histology and postoperative imaging. *Neurosurgery* 74:310-319; discussion 319-320. doi:10.1227/NEU.0000000000000267
24. Stummer W, Tonn JC, Mehdorn HM, Nestler U, Franz K, Goetz C, Bink A, Pichlmeier U (2011) Counterbalancing risks and gains from extended resections in malignant glioma surgery: a supplemental analysis from the randomized 5-aminolevulinic acid glioma resection study. *Clinical article. J Neurosurg* 114:613-623. doi:10.3171/2010.3.jns097
25. Stupp R, Mason WP, van den Bent MJ, Weller M, Fisher B, Taphoorn MJ, Belanger K, Brandes AA, Marosi C, Bogdahn U, Curschmann J, Janzer RC, Ludwin SK, Gorlia T, Allgeier A, Lacombe D, Cairncross JG, Eisenhauer E, Mirimanoff RO (2005) Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *The New England journal of medicine* 352:987-996. doi:10.1056/NEJMoa043330
26. Yamada S, Muragaki Y, Maruyama T, Komori T, Okada Y (2015) Role of neurochemical navigation with 5-aminolevulinic acid during intraoperative MRI-guided resection of intracranial malignant gliomas. *Clin Neurol Neurosurg* 130:134-139. doi:10.1016/j.clineuro.2015.01.005

Table 1. Patient and tumor characteristics		n=100
Gender	Male	52
	Female	48
Age	Range	26-80
	Mean	58,73 ±11,91
	Median	61
Pre-operative KPS	Range	20-100
	Median	90
	>70	84
	≤70	16
Tumor location	Eloquent	58
	Non-eloquent	42
Histology	GBM	90
	Anaplastic astrocytoma	3
	Anaplastic oligodendroglioma	1
	Anaplastic ependymoma	1
	Grade II astrocytoma	1
	Grade II oligodendroglioma	1
	Grade III meningioma	1
	Necrosis	1
	Vasculitis	1
Treatment after surgery: GBM	Stupp protocol	76
	Bevacizumab+irinotecan	1
	Palliative care	6
	Best supportive care	7

Table 2. Cross tabulation

		Resection			Total (n=90)	p
		Partial (n=23)	Subtotal (n=17)	Gross total (n=50)		
Sex	Male	13	5	29	47	0,112
	Female	10	12	21	43	
Age	<65	11	10	35	56	0,183
	≥65	12	7	15	34	
KPS pre-op	≤70	7	3	5	15	0,093
	>70	16	14	45	75	
Patients	1-45	13	9	23	45	0,680
	45-90	10	8	26	45	




FIG 1. Different extents of resection stratified by eloquent/non-eloquent area of tumor (p=0,019)




FIG. 2. Graphs showing Kaplan-Meier overall survival curves. A. Overall survival – 16,27 ± 1,39 months B. Survival stratified by age – <65 = 18,01±1,57 months; ≥65 = 13,22±2,31 months. C. Survival stratified by preoperative KPS score - KPS>70=17,36±1,54 months; KPS≤70=10,13±2,00 months. D. Survival stratified by tumor area (eloquent/non-eloquent) – eloquent = 13,71±1,32 months (median=12,00±1,55); non-eloquent = 19,32±2,41 months. E. Survival stratified by extent of resection – GTR>90%=17,52±1,68 months; PR = 13,130±2,62 months.




FIG. 3. Graphs showing Kaplan-Meier progression-free survival curves. A. Progression-free survival – $9,73 \pm 1,19$ months. B. Survival stratified by age – $<65=9,78 \pm 1,17$ months; $\geq 65=9,22 \pm 2,25$ months. C. Survival stratified by preoperative KPS score - $KPS > 70=10,48 \pm 1,37$ months; $KPS \leq 70=5,49 \pm 1,10$ months. D. Survival stratified by tumor area (eloquent/non-eloquent) – eloquent= $7,15 \pm 0,81$ months; non-eloquent= $13,58 \pm 2,59$ months. E. Survival stratified by extent of resection – $GTR > 90\%=11,33 \pm 1,54$ months; $PR=5,30 \pm 0,93$ months.



Acta Neurochirurgica

The European Journal of Neurosurgery

Editor-in-Chief: Tiit **Mathiesen**

ISSN: 0001-6268 (print version)

ISSN: 0942-0940 (electronic version)

Journal no. 701



Instructions for Authors

TYPES OF PAPERS

A checklist is provided to aid authors in preparing a manuscript that meets the standards of the journal. It is suggested that the completed checklist be uploaded during the submission process, but this is not required. The checklist is available [here](#)

Please state clearly the format of your paper, i.e. Original Article, Case Report, Letter to the Editor, Review Article, Technical Note.

Subdivision of the paper

Original Articles, Review Articles or Technical Notes.

For an original research paper the customary format consists of: abstract, introduction, methods and materials, results, discussion, references.

- The summary should be a short survey of the paper, structured using 4 headings: Background/ Method/ Results/ Conclusions. The abstract can run to a maximum of one full type-written page (300 words). Four to six keywords should be provided.
- The introduction should summarize the underlying purpose of the paper.
- Methods and materials must give short but precise descriptions, for instance of the kind of experiments and how they were done, the characteristics of patients or other humans participating as subjects, the nature of observations made and the specific approach to statistical considerations. This section should enable the reader to analyse, to repeat or to continue the work presented.
- Results must contain sufficient information to enable appreciation and

assessment of the original findings, e.g. absolute numbers in subgroups should be reported, preferably in tabular format, rather than only as a percentage. Where a normal distribution is assumed as a basis for summarizing data (e.g. as a mean), this should be justified and measures should be specified. Statistical tests should be presented in sufficient detail to convey the full result, not only as a “p” value.

- In discussion a systematic, structured approach is advised. A statement of the principal findings demonstrated by the work should be followed by a “self-critique”, focusing on the strengths and weaknesses of the study and how these may influence the validity and robustness of the findings. The relation of the work to other, relevant previous work, especially any differences, should be reviewed and analyzed concisely. Finally, the implications of the findings of the study for advancing knowledge and understanding, for neurosurgical practice and for further work should be presented.

Case reports.

Case reports should be limited to 6 pages of typescript (1500 words), including tables and illustrations. An abstract in a single paragraph, maximum 100 words, should be included. The presentation should both adequately characterise the features of the subject(s) reported and make clear what generalisable deductions can be drawn.

Letters to the Editor should be a maximum of 750 words and may include a table or figure and references. The format for submissions of editorials, reviews, personal communications, concept articles, biographies and historical articles can be varied as appropriate to the topic, consistent with clarity and brevity.

How I do It

This new section in the journal is for showing how surgical procedures are carried out.

Editor of this section is Prof. Pierre-Hugues Roche Please submit your paper following to the instructions in the download below. Visit the website below to check your connection speed prior to uploading:

The final categorisation of an accepted submission is the responsibility of the Editor in Chief.

How I do it - Instructions (pdf, 52 kB) Connection speed check

MANUSCRIPT SUBMISSION

Manuscript Submission

Submission of a manuscript implies: that the work described has not been published before; that it is not under consideration for publication anywhere else; that its publication has been approved by all co-authors, if any, as well as by the responsible authorities – tacitly or explicitly – at the institute where the work has been carried out. The publisher will not be held legally responsible should there be any claims for compensation.

Permissions

Authors wishing to include figures, tables, or text passages that have already been published elsewhere are required to obtain permission from the copyright owner(s) for both the print and online format and to include evidence that such permission has been granted when submitting their papers. Any material received without such evidence will be assumed to originate from the authors.

Online Submission

Please follow the hyperlink “Submit online” on the right and upload all of your manuscript files following the instructions given on the screen.

Please note:

During the submission process, it is possible to upload the completed “Checklist for Authors” as an additional file. This is recommended to ease the review process and aid the reviewers in suggesting revisions, but is not required. The “Checklist for Authors” is available [here](#)

TITLE PAGE

The word.doc file of your submission should be submitted as follows: Page 1: Title Page The title page should include:

Title of Paper Names(s) of all authors; Affiliation; Details of corresponding author-Full name, Affiliation, Full postal address, E-Mail/Fax details; Presentation at a conference Clinical; Trial Registration number if required Number the page-Title page will be page#1

Page 2: Abstract

A structured abstract is required between 150-250 words to be divided into

the following headings:

Background; Methods; Results; Conclusions

Keywords Please provide 4 to 6 keywords which can be used for indexing purposes

TEXT

[Text Formatting](#)

Manuscripts should be submitted in Word.

Use a normal, plain font (e.g., 10-point Times Roman) for text. Use italics for emphasis. Use the automatic page numbering function to number the pages. Do not use field functions.

Use tab stops or other commands for indents, not the space bar.

Use the table function, not spreadsheets, to make tables.

Use the equation editor or MathType for equations.

Save your file in docx format (Word 2007 or higher) or doc format (older Word versions).

Manuscripts with mathematical content can also be submitted in LaTeX.

LaTeX macro package (zip, 181 kB)

[Headings](#)

Please use no more than three levels of displayed headings.

[Abbreviations](#)

Abbreviations should be defined at first mention and used consistently thereafter.

[Footnotes](#)

Footnotes can be used to give additional information, which may include the citation of a reference included in the reference list. They should not consist solely of a reference citation, and they should never include the bibliographic details of a reference. They should also not contain any figures or tables.

Footnotes to the text are numbered consecutively; those to tables should be indicated by superscript lower-case letters (or asterisks for significance values and other statistical data). Footnotes to the title or the authors of the article are not given reference symbols.

Always use footnotes instead of endnotes.

Acknowledgments

Acknowledgments of people, grants, funds, etc. should be placed in a separate section on the title page. The names of funding organizations should be written in full.

REFERENCES

The list of references should only include works that are cited in the text and have been previously published or accepted for publication.

Personal communications should only be mentioned in the text.

In the text, references should be noted in square brackets eg: [1] and listed in alphabetical order in the reference list.

Papers which have been accepted for publication but not yet published should be included in the list of references with the name of the journal and the Digital Object Identifier (DOI) code of the cited literature. The author is responsible for the accuracy of the references.

Following some examples for the style of references: Journal articles

(1) Franzini A, Messina G, Leone M, Broggi G (2009) Occipital nerve stimulation (ONS). Surgical technique and prevention of late electrode migration. *Acta Neurochir* 151:861-865

(2) Pinsker MO, Volkmann J, Falk D, Herzog F, Steigerwald F, Deuschl G, Mehdorn HM (2009) Deep brain stimulation of the internal globus pallidus in dystonia: target localisation under general anaesthesia. *Acta Neurochir* 151:751-758

(3) Powell M (2009) Microscope and endoscopic pituitary surgery. *Acta Neurochir* 151:723-728 Article published on-line with DOI

(4) Helmy A, Timofeev I, Palmer C, Gore A, Menon DK, Hutchinson PJ (2010) Hierarchical log linear analysis of admission blood parameters and clinical outcome following traumatic brain injury *Acta Neurochir* DOI:

10.1007/S00701-009-0584-y

Books

Frontera J (2009) Decision making in neurocritical care Thieme, Stuttgart

Book Chapter

Hovind HJ (1986) Traumatic birth injuries. In: Raimondi AJ, Choux M, Di Rocco C (eds) Head injuries in the newborn and infant. (Principles of pediatric neurosurgery) Springer, Berlin Heidelberg New York, pp 87–109

Online Document

Human D, Fluss SS (2001) The world medical association's declaration of Helsinki: Historical and contemporary perspectives http://www.wma.net/e/ethicsunit/pdf/draft_historical_contemporary_perspectives.pdf. Accessed 8 Aug 2009

Always use the standard abbreviation of a journal's name according to the ISSN List of Title Word Abbreviations.

ISSN List of the Title Word Abbreviations

If you are unsure, please use the full journal title.

For authors using EndNote, Springer provides an output style that supports the formatting of in- text citations and reference list.

Endnote style (zip, 2 kB)

TABLES

All tables are to be numbered using Arabic numerals. Tables should always be cited in text in consecutive numerical order.

For each table, please supply a table caption (title) explaining the components of the table.

Identify any previously published material by giving the original source in the form of a reference at the end of the table caption.

Footnotes to tables should be indicated by superscript lower-case letters (or asterisks for significance values and other statistical data) and included beneath the table body.

ARTWORK AND ILLUSTRATIONS GUIDELINES

Electronic Figure Submission

Supply all figures electronically. Indicate what graphics program was used to create the artwork.

For vector graphics, the preferred format is EPS; for halftones, please use TIFF format. MSOffice files are also acceptable.


Vector graphics containing fonts must have the fonts embedded in the files. Name your figure files with "Fig" and the figure number, e.g., Fig1.eps.

Line Art

Definition: Black and white graphic with no shading.

Do not use faint lines and/or lettering and check that all lines and lettering within the figures are legible at final size.

All lines should be at least 0.1 mm (0.3 pt) wide.




Scanned line drawings and line drawings in bitmap format should have a minimum resolution of 1200 dpi.

Vector graphics containing fonts must have the fonts embedded in the files.

Halftone Art


Definition: Photographs, drawings, or paintings with fine shading, etc. If any magnification is used in the photographs, indicate this by using scale bars within the figures themselves.

Halftones should have a minimum resolution of 300 dpi.



⌘ ⌘

Combination Art



Definition: a combination of halftone and line art, e.g., halftones containing line drawing, extensive lettering, color diagrams, etc. Combination artwork

should have a minimum resolution of 600 dpi.

Color Art

Color art is free of charge for online publication. If black and white will be shown in the print version, make sure that the main information will still be visible. Many colors are not distinguishable from one another when converted to black and white. A simple way to check this is to make a xerographic copy to see if the necessary distinctions between the different colors are still apparent. If the figures will be printed in black and white, do not refer to color in the captions. Color illustrations should be submitted as RGB (8 bits per channel).

Figure Lettering

To add lettering, it is best to use Helvetica or Arial (sans serif fonts).

Keep lettering consistently sized throughout your final-sized artwork, usually about 2–3 mm (8–12 pt).

Variance of type size within an illustration should be minimal, e.g., do not use 8-pt type on an axis and 20-pt type for the axis label.

Avoid effects such as shading, outline letters, etc. Do not include titles or captions within your illustrations.

Figure Numbering

All figures are to be numbered using Arabic numerals. Figures should always be cited in text in consecutive numerical order. Figure parts should be denoted by lowercase letters (a, b, c, etc.). If an appendix appears in your article and it contains one or more figures, continue the consecutive numbering of the main text. Do not number the appendix figures,

"A1, A2, A3, etc." Figures in online appendices (Electronic Supplementary Material) should, however, be numbered separately.

Figure Captions

Each figure should have a concise caption describing accurately what the figure depicts. Include the captions in the text file of the manuscript, not in the figure file.

Figure captions begin with the term **Fig.** in bold type, followed by the figure number, also in bold type.

No punctuation is to be included after the number, nor is any punctuation to be placed at the end of the caption.

Identify all elements found in the figure in the figure caption; and use boxes, circles, etc., as coordinate points in graphs.

Identify previously published material by giving the original source in the form of a reference citation at the end of the figure caption.

Figure Placement and Size

Figures should be submitted separately from the text, if possible. When preparing your figures, size figures to fit in the column width. For most journals the figures should be 39 mm, 84 mm, 129 mm, or 174 mm wide and not higher than 234 mm. For books and book-sized journals, the figures should be 80 mm or 122 mm wide and not higher than 198 mm.

Permissions

If you include figures that have already been published elsewhere, you must obtain permission from the copyright owner(s) for both the print and online format. Please be aware that some publishers do not grant electronic rights for free and that Springer will not be able to refund any costs that may have occurred to receive these permissions. In such cases, material from other sources should be used.

Accessibility

In order to give people of all abilities and disabilities access to the content of your figures, please make sure that

All figures have descriptive captions (blind users could then use a text-to-speech software or a text-to-Braille hardware) Patterns are used instead of or in addition to colors for conveying information (colorblind users would then be able to distinguish the visual elements)

Any figure lettering has a contrast ratio of at least 4.5:1

ELECTRONIC SUPPLEMENTARY MATERIAL

Springer accepts electronic multimedia files (animations, movies, audio, etc.) and other supplementary files to be published online along with an article or a book chapter. This feature can add dimension to the author's article, as certain information cannot be printed or is more convenient in electronic form.

Before submitting research datasets as electronic supplementary material, authors should read the journal's Research data policy. We encourage research data to be archived in data repositories wherever possible.

Submission

Supply all supplementary material in standard file formats. Please include in each file the following information: article title, journal name, author names; affiliation and e-mail address of the corresponding author. To accommodate user downloads, please keep in mind that larger-sized files may require very long download times and that some users may experience other problems during downloading.

Audio, Video, and Animations

Aspect ratio: 16:9 or 4:3 Maximum file size: 25 GB Minimum video duration: 1 sec Supported file formats: avi, wmv, mp4, mov, m2p, mp2, mpg, mpeg, flv, mxf, mts, m4v, 3gp

Text and Presentations

Submit your material in PDF format; .doc or .ppt files are not suitable for long-term viability. A collection of figures may also be combined in a PDF file.

Spreadsheets

Spreadsheets should be submitted as .csv or .xlsx files (MS Excel).

Specialized Formats

Specialized format such as .pdb (chemical), .wrl (VRML), .nb (Mathematica notebook), and .tex can also be supplied.

Collecting Multiple Files

It is possible to collect multiple files in a .zip or .gz file.

Numbering

If supplying any supplementary material, the text must make specific mention of the material as a citation, similar to that of figures and tables. Refer to the supplementary files as "Online Resource", e.g., "... as shown in the animation (Online Resource 3)", "... additional data are given in Online Resource 4". Name the files consecutively, e.g. "ESM_3.mpg",

“ESM_4.pdf”.

Captions

For each supplementary material, please supply a concise caption describing the content of the file.

Processing of supplementary files

Electronic supplementary material will be published as received from the author without any conversion, editing, or reformatting.

Accessibility

In order to give people of all abilities and disabilities access to the content of your supplementary files, please make sure that

The manuscript contains a descriptive caption for each supplementary material. Video files do not contain anything that flashes more than three times per second (so that users prone to seizures caused by such effects are not put at risk).

Audio, Video, and Animations

Use the file type „supplementary files“ to upload the video. The time for uploading cannot be longer than 15 mins. If the files are large, break down the video to 2 or 3 parts and upload separately. A fast internet connection may be required.

INTEGRITY OF RESEARCH AND REPORTING

Ethical standards

Manuscripts submitted for publication must contain a statement to the effect that all human and animal studies have been approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki and its later amendments.

It should also be stated clearly in the text that all persons gave their informed consent prior to their inclusion in the study. Details that might disclose the identity of the subjects under study should be omitted.

These statements should be added in a separate section before the reference list. If these statements are not applicable, authors should state:

The manuscript does not contain clinical studies or patient data.

The editors reserve the right to reject manuscripts that do not comply with the above-mentioned requirements. The author will be held responsible for false statements or failure to fulfill the above-mentioned requirements

Conflict of interest

Authors must indicate whether or not they have a financial relationship with the organization that sponsored the research. This note should be added in a separate section before the reference list.

If no conflict exists, authors should state: The authors declare that they have no conflict of interest.

ETHICAL RESPONSIBILITIES OF AUTHORS

This journal is committed to upholding the integrity of the scientific record. As a member of the Committee on Publication Ethics (COPE) the journal will follow the COPE guidelines on how to deal with potential acts of misconduct.

Authors should refrain from misrepresenting research results which could damage the trust in the journal, the professionalism of scientific authorship, and ultimately the entire scientific endeavour. Maintaining integrity of the research and its presentation can be achieved by following the rules of good scientific practice, which include:

The manuscript has not been submitted to more than one journal for simultaneous consideration.

The manuscript has not been published previously (partly or in full), unless the new work concerns an expansion of previous work (please provide transparency on the re-use of material to avoid the hint of text-recycling (“self-plagiarism”)).

A single study is not split up into several parts to increase the quantity of submissions and submitted to various journals or to one journal over time (e.g. “salami-publishing”).

No data have been fabricated or manipulated (including images) to support your conclusions

No data, text, or theories by others are presented as if they were the author’s own (“plagiarism”). Proper acknowledgements to other works

must be given (this includes material that is closely copied (near verbatim), summarized and/or paraphrased), quotation marks are used for verbatim copying of material, and permissions are secured for material that is copyrighted.

Important note: the journal may use software to screen for plagiarism.

Consent to submit has been received explicitly from all co-authors, as well as from the responsible authorities - tacitly or explicitly - at the institute/organization where the work has been carried out, **before** the work is submitted.

Authors whose names appear on the submission have contributed sufficiently to the scientific work and therefore share collective responsibility and accountability for the results.

Authors are strongly advised to ensure the correct author group, corresponding author, and order of authors at submission. Changes of authorship or in the order of authors are **not** accepted **after** acceptance of a manuscript.

Adding and/or deleting authors and/or changing the order of authors **at revision stage** may be justifiably warranted. A letter must accompany the revised manuscript to explain the reason for the change(s) and the contribution role(s) of the added and/or deleted author(s). Further documentation may be required to support your request.

Requests for addition or removal of authors as a result of authorship disputes after acceptance are honored after formal notification by the institute or independent body and/or when there is agreement between all authors.

Upon request authors should be prepared to send relevant documentation or data in order to verify the validity of the results. This could be in the form of raw data, samples, records, etc. Sensitive information in the form of confidential proprietary data is excluded.

If there is a suspicion of misconduct, the journal will carry out an investigation following the COPE guidelines. If, after investigation, the allegation seems to raise valid concerns, the accused author will be contacted and given an opportunity to address the issue. If misconduct has been established beyond reasonable doubt, this may result in the Editor-in-Chief's implementation of the following measures, including, but not limited to:

If the article is still under consideration, it may be rejected and returned to the author. If the article has already been published online, depending on the nature and severity of the infraction, either an erratum will be placed with the article or in severe cases complete retraction of the article will occur. The reason must be given in the published erratum or retraction note. Please note that retraction means that the paper is **maintained on the platform**, watermarked "retracted" and explanation for the retraction is provided in a note linked to the watermarked article.

The author's institution may be informed.

COMPLIANCE WITH ETHICAL STANDARDS

To ensure objectivity and transparency in research and to ensure that accepted principles of ethical and professional conduct have been followed, authors should include information regarding sources of funding, potential conflicts of interest (financial or non-financial), informed consent if the research involved human participants, and a statement on welfare of animals if the research involved animals.

Authors should include the following statements (if applicable) in a separate section entitled "Compliance with Ethical Standards" when submitting a paper:

Disclosure of potential conflicts of interest Research involving Human Participants and/or Animals Informed consent

Please note that standards could vary slightly per journal dependent on their peer review policies (i.e. single or double blind peer review) as well as per journal subject discipline. Before submitting your article check the instructions following this section carefully.

The corresponding author should be prepared to collect documentation of compliance with ethical standards and send if requested during peer review or after publication.

The Editors reserve the right to reject manuscripts that do not comply with the above-mentioned guidelines. The author will be held responsible for false statements or failure to fulfill the above mentioned guidelines.

DISCLOSURE OF POTENTIAL CONFLICTS OF INTEREST

Authors must disclose all relationships or interests that could have direct or potential influence or impart bias on the work. Although an author may not feel there is any conflict, disclosure of relationships and interests provides

a more complete and transparent process, leading to an accurate and objective assessment of the work. Awareness of a real or perceived conflicts of interest is a perspective to which the readers are entitled. This is not meant to imply that a financial relationship with an organization that sponsored the research or compensation received for consultancy work is inappropriate. Examples of potential conflicts of interests **that are directly or indirectly related to the research** may include but are not limited to the following:

Research grants from funding agencies (please give the research funder and the grant number)

Honoraria for speaking at symposia

Financial support for attending symposia

Financial support for educational programs

Employment or consultation

Support from a project sponsor

Position on advisory board or board of directors or other type of management relationships

Multiple affiliations Financial relationships, for example equity ownership or investment interest Intellectual property rights (e.g. patents, copyrights and royalties from such rights) Holdings of spouse and/or children that may have financial interest in the work

In addition, interests that go beyond financial interests and compensation (non-financial interests) that may be important to readers should be disclosed. These may include but are not limited to personal relationships or competing interests directly or indirectly tied to this research, or professional interests or personal beliefs that may influence your research.

The corresponding author collects the conflict of interest disclosure forms from all authors. In author collaborations where formal agreements for representation allow it, it is sufficient for the corresponding author to sign the disclosure form on behalf of all authors. Examples of forms can be found

here:

The corresponding author will include a summary statement in the text of

the manuscript in a separate section before the reference list, that reflects what is recorded in the potential conflict of interest disclosure form(s).

Please make sure to submit all Conflict of Interest disclosure forms together with the manuscript.

See below examples of disclosures: **Funding:** This study was funded by X (grant number X).

Conflict of Interest: Author A has received research grants from Company A. Author B has received a speaker honorarium from Company X and owns stock in Company Y. Author C is a member of committee Z.

If no conflict exists, the authors should state: Conflict of Interest: The authors declare that they have no conflict of interest.

RESEARCH INVOLVING HUMAN PARTICIPANTS AND/OR ANIMALS

1) Statement of human rights

When reporting studies that involve human participants, authors should include a statement that the studies have been approved by the appropriate institutional and/or national research ethics committee and have been performed in accordance with the ethical standards as laid down in the 1964 Declaration of Helsinki and its later amendments or comparable ethical standards.

If doubt exists whether the research was conducted in accordance with the 1964 Helsinki Declaration or comparable standards, the authors must explain the reasons for their approach, and demonstrate that the independent ethics committee or institutional review board explicitly approved the doubtful aspects of the study.

The following statements should be included in the text before the References section:

Ethical approval: “All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.”

For retrospective studies, please add the following sentence: “For this type of study formal consent is not required.”

2) Statement on the welfare of animals

The welfare of animals used for research must be respected. When reporting experiments on animals, authors should indicate whether the international, national, and/or institutional guidelines for the care and use of animals have been followed, and that the studies have been approved by a research ethics committee at the institution or practice at which the studies were conducted (where such a committee exists).

For studies with animals, the following statement should be included in the text before the References section:

Ethical approval: “All applicable international, national, and/or institutional guidelines for the care and use of animals were followed.”

If applicable (where such a committee exists): “All procedures performed in studies involving animals were in accordance with the ethical standards of the institution or practice at which the studies were conducted.”

If articles do not contain studies with human participants or animals by any of the authors, please select one of the following statements:

“This article does not contain any studies with human participants performed by any of the authors.”

“This article does not contain any studies with animals performed by any of the authors.”

“This article does not contain any studies with human participants or animals performed by any of the authors.”

INFORMED CONSENT

All individuals have individual rights that are not to be infringed. Individual participants in studies have, for example, the right to decide what happens to the (identifiable) personal data gathered, to what they have said during a study or an interview, as well as to any photograph that was taken. Hence it is important that all participants gave their informed consent in writing prior to inclusion in the study. Identifying details (names, dates of birth, identity numbers and other information) of the participants that were studied should not be published in written descriptions, photographs, and genetic profiles unless the information is essential for scientific purposes and the participant (or parent or guardian if the participant is incapable) gave written informed consent for publication. Complete anonymity is difficult to achieve in some cases, and informed consent should be obtained if there is any doubt. For example, masking the eye region in photographs of participants is inadequate protection of anonymity. If

identifying characteristics are altered to protect anonymity, such as in genetic profiles, authors should provide assurance that alterations do not distort scientific meaning.

The following statement should be included:

Informed consent: “Informed consent was obtained from all individual participants included in the study.”

If identifying information about participants is available in the article, the following statement should be included:

“Additional informed consent was obtained from all individual participants for whom identifying information is included in this article.”

ENGLISH LANGUAGE EDITING

For editors and reviewers to accurately assess the work presented in your manuscript you need to ensure the English language is of sufficient quality to be understood. If you need help with writing in English you should consider:

Asking a colleague who is a native English speaker to review your manuscript for clarity. Visiting the English language tutorial which covers the common mistakes when writing in English.

Using a professional language editing service where editors will improve the English to ensure that your meaning is clear and identify problems that require your review. Two such services are provided by our affiliates Nature Research Editing Service and American Journal Experts. Springer authors are entitled to a 10% discount on their first submission to either of these services, simply follow the links below.

English language tutorial [Nature Research Editing Service](#) [American Journal Experts](#)

Please note that the use of a language editing service is not a requirement for publication in this journal and does not imply or guarantee that the article will be selected for peer review or accepted.

If your manuscript is accepted it will be checked by our copyeditors for spelling and formal style before publication.

RESEARCH DATA POLICY

The journal encourages authors, where possible and applicable, to deposit

data that support the findings of their research in a public repository. Authors and editors who do not have a preferred repository should consult Springer Nature's list of repositories and research data policy.

List of Repositories Research Data Policy

General repositories - for all types of research data - such as figshare and Dryad may also be used.

Datasets that are assigned digital object identifiers (DOIs) by a data repository may be cited in the reference list. Data citations should include the minimum information recommended by DataCite: authors, title, publisher (repository name), identifier.

DataCite

Springer Nature provides a research data policy support service for authors and editors, which can be contacted at **researchdata@springernature.com**.

This service provides advice on research data policy compliance and on finding research data repositories. It is independent of journal, book and conference proceedings editorial offices and does not advise on specific manuscripts.

Helpdesk

AFTER ACCEPTANCE

Upon acceptance of your article you will receive a link to the special Author Query Application at Springer's web page where you can sign the Copyright Transfer Statement online and indicate whether you wish to order OpenChoice and offprints.

Once the Author Query Application has been completed, your article will be processed and you will receive the proofs.

[Copyright transfer](#)

Authors will be asked to transfer copyright of the article to the Publisher (or grant the Publisher exclusive publication and dissemination rights). This will ensure the widest possible protection and dissemination of information under copyright laws.

[Offprints](#)

Offprints can be ordered by the corresponding author.

Color illustrations

Publication of color illustrations is free of charge.

Proof reading

The purpose of the proof is to check for typesetting or conversion errors and the completeness and accuracy of the text, tables and figures. Substantial changes in content, e.g., new results, corrected values, title and authorship, are not allowed without the approval of the Editor.

After online publication, further changes can only be made in the form of an Erratum, which will be hyperlinked to the article.

Online First

The article will be published online after receipt of the corrected proofs. This is the official first publication citable with the DOI. After release of the printed version, the paper can also be cited by issue and page numbers.

OPEN CHOICE

In addition to the normal publication process (whereby an article is submitted to the journal and access to that article is granted to customers who have purchased a subscription), Springer provides an alternative publishing option: Springer Open Choice. A Springer Open Choice article receives all the benefits of a regular subscription-based article, but in addition is made available publicly through Springer's online platform SpringerLink.

Open Choice

Copyright and license term – CC BY

Open Choice articles do not require transfer of copyright as the copyright remains with the author. In opting for open access, the author(s) agree to publish the article under the Creative Commons Attribution License.

Find more about the license agreement

EDITORIAL OFFICE

D. Epstein Di-Ep Biomedical Editorial Services 10 Sandyford Place Glasgow G3-7NB Scotland, UK Fax +44-141-6169095 e-mail:

diana.epstein@di-ep.com